arXiv:1205.5987v1 [math.CA] 27 May 2012

HERMITE-HADAMARD TYPE INEQUALITIES FOR FUNCTIONS WHOSE DERIVATIVES ARE STRONGLY $\varphi-$CONVEX

İMDAT İŞCAN AND ERDAL UNLUYOL

Abstract. In this paper several inequalities of the right-hand side of Hermite-Hadamard's inequality are obtained for the class of functions whose derivatives in absolutely value at certain powers are strongly φ-convex with modulus c>0.

1. Introduction

Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$, then

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_{a}^{b} f(x)dx \leq \frac{f(a)+f(b)}{2}. \tag{1.1}$$

This doubly inequality is known in the literature as Hermite-Hadamard integral inequality for convex functions. This inequality plays an important role in convex analysis and it has a huge literature dealing with its applications, various generalizations and refinements (see [1, 2, 3, 4, 5, 8, 11]).

Let us consider a function $\varphi : [a, b] \longrightarrow [a, b]$, where $[a, b] \subset \mathbb{R}$.Youness have defined the $\varphi-$convex functions in [12] :

Definition 1. *A function $f : [a, b]\rightarrow \mathbb{R}$ is said to be $\varphi-$convex on $[a, b]$ if for every two points $x, y \in [a, b]$ and $t \in [0, 1]$ the following inequality holds:*

$$f\left(t\varphi(x) + (1-t)\varphi(y)\right) \leq tf\left(\varphi(x)\right) + (1-t)f\left(\varphi(y)\right).$$

Obviously, if function φ is the identity, then the classical convexity is obtained from the previous definition.

Recall also that a function $f : I\rightarrow \mathbb{R}$ is called strongly convex with modulus $c > 0$, if

$$f\left(tx + (1-t)y\right) \leq tf\left(x\right) + (1-t)f\left(y\right) - ct(1-t)\left(x-y\right)^2$$

for all $x, y \in I$ and $t \in (0, 1)$. Strongly convex functions have been introduced by Polyak in [9] and they play an important role in optimization theory and mathematical economics. Various properties and applications of them can be found in the literature see [6, 7, 9].

Date: May 26, 2012.

1991 *Mathematics Subject Classification.* 26D10, 26A51.

Key words and phrases. Hermite-Hadamard's inequalities, φ-convex functions, strongly φ-convex with modulus c>0,

Let $(X, \|.\|)$ be a real normed space, D stands for a convex subset of X, $\varphi : D \longrightarrow D$ is a given function and c is a positive constant. In [10] Sarikaya have introduced the notion of the strongly $\varphi-$convex functions with modulus c and some properties of them. Moreover in his paper, Sarikaya have presented a version Hermite-Hadamard-type inequalities for strongly $\varphi-$convex functions as follows:

Definition 2. *A function $f : D\to \mathbb{R}$ is said to be strongly $\varphi-$convex with modulus c if*

$$f\left(t\varphi(x)+(1-t)\varphi(y)\right)\leq tf\left(\varphi(x)\right)+(1-t)f\left(\varphi(y)\right)-ct(1-t)\left\|\varphi(x)-\varphi(y)\right\|^{2}$$

for all $x, y \in D$ and $t \in [0, 1]$.

The notion of $\varphi-$convex function corresponds to the case $c = 0$.If function φ is the identity, then the strongly convexity with modulus $c > 0$ is obtained from the previous definition

Theorem 1. *If $f : [a, b]\to \mathbb{R}$ is strongly $\varphi-$convex with modulus $c > 0$ for the continuous function $\varphi : [a, b] \longrightarrow [a, b]$, then*

$$\begin{aligned} f\left(\frac{\varphi(a)+\varphi(b)}{2}\right)+\frac{c}{12}\left(\varphi(b)-\varphi(a)\right)^{2} &\leq \frac{1}{\varphi(b)-\varphi(a)}\int\limits_{\varphi(a)}^{\varphi(b)}f(x)dx \\ &\leq \frac{f\left(\varphi(a)\right)+f\left(\varphi(b)\right)}{2}-\frac{c}{6}\left(\varphi(b)-\varphi(a)\right)^{2} \end{aligned}$$

The main aim of this paper is to establish new inequalities of Hermite-Hadamard type for the class of functions whose derivatives at certain powers are strongly $\varphi-$convex with modulus c.

2. Inequalities for functions whose derivatives are strongly $\varphi-$convex with modulus c

In order to prove our main resuls we need the following lemma:

Lemma 1. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I°, $a, b \in I$ with $a < b$. If $f' \in L[a, b]$ and $\varphi : [a, b] \longrightarrow [a, b]$ a continuous function with $\varphi(a) < \varphi(b)$. then the following equality holds:*

(2.1)
$$\frac{f\left(\varphi(a)\right)+f\left(\varphi(b)\right)}{2}-\frac{1}{\varphi(b)-\varphi(a)}\int\limits_{\varphi(a)}^{\varphi(b)}f(x)dx=\frac{\varphi(b)-\varphi(a)}{2}\int\limits_{0}^{1}\left(2t-1\right)f'\left(t\varphi(b)+(1-t)\varphi(a)\right)dt$$

Proof. By using partial integration in right hand of (2.1) equality, the proof is obvious. □

The next theorem gives a new the upper Hermite-Hadamard inequality for strongly $\varphi-$convex functions with modulus c as follows:

Theorem 2. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I°, $a, b \in I$ with $a < b$ such that $f' \in L[a, b]$. If $|f'|^q$ is strongly $\varphi-$convex functions with modulus c* *on* $[a,b]$ *for the continuous function* $\varphi:[a,b]\longrightarrow[a,b]$ *with* $\varphi(a)<\varphi(b)$ *and* $q\geq 1$ *then*

$$\begin{aligned}(2.2)\qquad &\left|\frac{f\left(\varphi(a)\right)+f\left(\varphi(b)\right)}{2}-\frac{1}{\varphi(b)-\varphi(a)}\int\limits_{\varphi(a)}^{\varphi(b)}f(x)dx\right| \\ \leq\ & \frac{\varphi(b)-\varphi(a)}{4}\left(\frac{\left|f'(\varphi(b))\right|^{q}+\left|f'(\varphi(a))\right|^{q}}{2}-\frac{c}{8}\left(\varphi(b)-\varphi(a)\right)^{2}\right)^{\frac{1}{q}}\end{aligned}$$

Proof. Suppose that $q=1$. From Lemma 1 and using the strongly $\varphi-$convexity functions with modulus c of $|f'|$, we have

$$\begin{aligned}&\left|\frac{f\left(\varphi(a)\right)+f\left(\varphi(b)\right)}{2}-\frac{1}{\varphi(b)-\varphi(a)}\int\limits_{\varphi(a)}^{\varphi(b)}f(x)dx\right| \\ \leq\ & \frac{\varphi(b)-\varphi(a)}{2}\int\limits_{0}^{1}\left|2t-1\right|\left|f'(t\varphi(b)+(1-t)\varphi(a))\right|dt \\ \leq\ & \frac{\varphi(b)-\varphi(a)}{2}\int\limits_{0}^{1}\left|2t-1\right|\left[t\left|f'(\varphi(b))\right|+(1-t)\left|f'(\varphi(a))\right|-ct\left(1-t\right)\left(\varphi(b)-\varphi(a)\right)^{2}\right]dt\end{aligned}$$

We have

$$\int\limits_{0}^{1}\left|2t-1\right|tdt=\frac{1}{4},\ \int\limits_{0}^{1}\left|2t-1\right|(1-t)dt=\frac{1}{4}$$

and

$$\int\limits_{0}^{1}\left|2t-1\right|t\left(1-t\right)dt=\frac{1}{16}$$

hence we obtain

$$\begin{aligned}&\left|\frac{f\left(\varphi(a)\right)+f\left(\varphi(b)\right)}{2}-\frac{1}{\varphi(b)-\varphi(a)}\int\limits_{\varphi(a)}^{\varphi(b)}f(x)dx\right| \\ \leq\ & \frac{\varphi(b)-\varphi(a)}{4}\left(\frac{\left|f'(\varphi(b))\right|+\left|f'(\varphi(a))\right|}{2}-\frac{c}{8}\left(\varphi(b)-\varphi(a)\right)^{2}\right)\end{aligned}$$

which copmletes the proof for this case.

Suppose now that $q \in (1,\infty)$. From Lemma 1 and using the Hölder's integral inequality, we have

$$\int_0^1 |2t-1| \left|f'(t\varphi(b)+(1-t)\varphi(a))\right| dt \tag{2.3}$$
$$\leq \frac{\varphi(b)-\varphi(a)}{2}\left(\int_0^1 |2t-1|\,dt\right)^{\frac{q-1}{q}}\left(\int_0^1 |2t-1|\left|f'(t\varphi(b)+(1-t)\varphi(a))\right|^q dt\right)^{\frac{1}{q}}$$

Since $|f'|^q$ is strongly $\varphi-$convex functions with modulus c on $[a,b]$, we know that for every $t \in [0,1]$

(2.4)

$$|f'(t\varphi(b)+(1-t)\varphi(a))|^q \leq t\,|f'(\varphi(b))|^q+(1-t)\,|f'(\varphi(a))|^q-ct\,(1-t)\,(\varphi(b)-\varphi(a))^2 .$$

From 2.1, 2.3 and 2.4, we have

$$\left|\frac{f\left(\varphi(a)\right)+f\left(\varphi(b)\right)}{2}-\frac{1}{\varphi(b)-\varphi(a)}\int_{\varphi(a)}^{\varphi(b)} f(x)dx\right|$$
$$\leq \frac{\varphi(b)-\varphi(a)}{4}\left(\frac{|f'(\varphi(b))|^q+|f'(\varphi(a))|^q}{2}-\frac{c}{8}\left(\varphi(b)-\varphi(a)\right)^2\right)^{\frac{1}{q}}$$

which completes the proof. □

Corollary 1. *Suppose that all the assumptions of Theorem2 are satisfied, in this case:*

(1) *For $c=0$, i.e. if $|f'|^q$ is $\varphi-$convex functions, we have*

$$\left|\frac{f\left(\varphi(a)\right)+f\left(\varphi(b)\right)}{2}-\frac{1}{\varphi(b)-\varphi(a)}\int_{\varphi(a)}^{\varphi(b)} f(x)dx\right| \tag{2.5}$$
$$\leq \frac{\varphi(b)-\varphi(a)}{4}\left(\frac{|f'(\varphi(b))|^q+|f'(\varphi(a))|^q}{2}\right)^{\frac{1}{q}}$$

(2) *For $\varphi(t)=t$ in (2.5) inequality :*

$$\left|\frac{f\left(a\right)+f\left(b\right)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right|$$
$$\leq \frac{b-a}{4}\left(\frac{|f'(b)|^q+|f'(a)|^q}{2}\right)^{\frac{1}{q}}$$

we get the same result in [8, Theorem 1].

(3) *If we take $\varphi(t) = t$ in (2.2) for strongly convex functions with modulus $c > 0$ on $[a, b]$ inequality, then we obtain*

$$\begin{aligned} &\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x)dx \right| \\ \leq\; & \frac{b-a}{4} \left(\frac{|f'(b)|^q + |f'(a)|^q}{2} - \frac{c}{8}(b-a)^2 \right)^{\frac{1}{q}} \end{aligned}$$

Theorem 3. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I°, $a, b \in I$ with $a < b$ such that $f' \in L[a, b]$. If $|f'|^q$ is strongly $\varphi-$convex functions with modulus c on $[a, b]$ for the continuous function $\varphi : [a, b] \longrightarrow [a, b]$ with $\varphi(a) < \frac{\varphi(a)+\varphi(b)}{2} < \varphi(b)$ and $q > 1$ then*

$$\begin{aligned} &\left| \frac{f(\varphi(a)) + f(\varphi(b))}{2} - \frac{1}{\varphi(b) - \varphi(a)} \int_{\varphi(a)}^{\varphi(b)} f(x)dx \right| \leq \frac{\varphi(b) - \varphi(a)}{4} \left(\frac{1}{(p+1)} \right)^{\frac{1}{p}} \\ &\times \left(\frac{1}{2}\right)^{\frac{1}{q}} \left[\begin{array}{c} \left(\left| f'\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \right|^q + |f'(\varphi(a))|^q - \frac{c}{3}(\varphi(b) - \varphi(a))^2 \right)^{\frac{1}{q}} \\ + \left(\left| f'\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \right|^q + |f'(\varphi(b))|^q - \frac{c}{3}(\varphi(b) - \varphi(a))^2 \right)^{\frac{1}{q}} \end{array} \right], \end{aligned} \tag{2.6}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From Lemma 1 and using the Hölder inequality, we have

$$\begin{aligned} &\left| \frac{f(\varphi(a)) + f(\varphi(b))}{2} - \frac{1}{\varphi(b) - \varphi(a)} \int_{\varphi(a)}^{\varphi(b)} f(x)dx \right| \\ \leq\; & \frac{\varphi(b) - \varphi(a)}{2} \left(\int_0^{\frac{1}{2}} (1-2t)^p \, dt \right)^{\frac{1}{p}} \left(\int_0^{\frac{1}{2}} |f'(t\varphi(b) + (1-t)\varphi(a))|^q \, dt \right)^{\frac{1}{q}} \\ &+ \frac{\varphi(b) - \varphi(a)}{2} \left(\int_{\frac{1}{2}}^1 (2t-1)^p \, dt \right)^{\frac{1}{p}} \left(\int_{\frac{1}{2}}^1 |f'(t\varphi(b) + (1-t)\varphi(a))|^q \, dt \right)^{\frac{1}{q}} \\ \leq\; & \frac{\varphi(b) - \varphi(a)}{2} \left(\frac{1}{2(p+1)} \right)^{\frac{1}{p}} \left(\frac{1}{4}\right)^{\frac{1}{q}} \left[\begin{array}{c} \left(\left| f'\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \right|^q + |f'(\varphi(a))|^q - \frac{c}{3}(\varphi(b) - \varphi(a))^2 \right)^{\frac{1}{q}} \\ + \left(\left| f'\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \right|^q + |f'(\varphi(b))|^q - \frac{c}{3}(\varphi(b) - \varphi(a))^2 \right)^{\frac{1}{q}} \end{array} \right] \end{aligned}$$

where we use the fact that

$$\int_0^{\frac{1}{2}} (1-2t)^p \, dt = \int_{\frac{1}{2}}^1 (2t-1)^p \, dt = \frac{1}{2(p+1)}$$

and by Theorem1 we get

$$\begin{aligned}
\int_{0}^{\frac{1}{2}} \left|f'(t\varphi(b)+(1-t)\varphi(a))\right|^{q} dt &= \frac{1}{2}\frac{2}{\varphi(b)-\varphi(a)} \int_{\varphi(a)}^{\frac{\varphi(a)+\varphi(b)}{2}} \left|f'(x)\right|^{q} \ dx \\
&\leq \left(\frac{1}{4}\right)\left(\left|f'\left(\frac{\varphi(a)+\varphi(b)}{2}\right)\right|^{q}+\left|f'\left(\varphi(a)\right)\right|^{q}-\frac{c}{3}\left(\varphi(b)-\varphi(a)\right)^{2}\right), \\
\frac{\lambda+\mu}{\mu}\int_{\frac{\lambda}{\lambda+\mu}}^{1} \left|f'(tb+(1-t)a)\right|^{q} \ dt &= \frac{1}{2}\frac{2}{\varphi(b)-\varphi(a)} \int_{\frac{\varphi(a)+\varphi(b)}{2}}^{\varphi(b)} \left|f'(x)\right|^{q} \ dx \\
&\leq \left(\frac{1}{4}\right)\left(\left|f'\left(\frac{\varphi(a)+\varphi(b)}{2}\right)\right|^{q}+\left|f'\left(\varphi(b)\right)\right|^{q}-\frac{c}{3}\left(\varphi(b)-\varphi(a)\right)^{2}\right).
\end{aligned}$$

□

Corollary 2. *Suppose that all the assumptions of Theorem3 are satisfied, in this case:*

1. *Since $|f'|^q$ is strongly $\varphi-$convex functions with modulus c, from (2.6) inequality we get*

$$\begin{aligned}
&\left|\frac{f\left(\varphi(a)\right)+f\left(\varphi(b)\right)}{2}-\frac{1}{\varphi(b)-\varphi(a)}\int_{\varphi(a)}^{\varphi(b)} f(x)dx\right| \\
\leq\ & \frac{\varphi(b)-\varphi(a)}{4}\left(\frac{1}{(p+1)}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1}{q}}\left[\begin{array}{c} \left(\frac{\left|f'(\varphi(b))\right|^{q}+3\left|f'(\varphi(a))\right|^{q}}{2}-\frac{7c}{12}\left(\varphi(b)-\varphi(a)\right)^{2}\right)^{\frac{1}{q}} \\ +\left(\frac{3\left|f'(\varphi(b))\right|^{q}+\left|f'(\varphi(a))\right|^{q}}{2}-\frac{7c}{12}\left(\varphi(b)-\varphi(a)\right)^{2}\right)^{\frac{1}{q}} \end{array}\right]
\end{aligned}$$

2. *For $c=0$, i.e. if $|f'|^q$ is $\varphi-$convex functions, we have*

$$\begin{aligned}
&\left|\frac{f\left(\varphi(a)\right)+f\left(\varphi(b)\right)}{2}-\frac{1}{\varphi(b)-\varphi(a)}\int_{\varphi(a)}^{\varphi(b)} f(x)dx\right| \\
\leq\ & \frac{\varphi(b)-\varphi(a)}{4}\left(\frac{1}{(p+1)}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1}{q}}\left[\begin{array}{c} \left(\left|f'\left(\frac{\varphi(a)+\varphi(b)}{2}\right)\right|^{q}+\left|f'\left(\varphi(a)\right)\right|^{q}\right)^{\frac{1}{q}} \\ +\left(\left|f'\left(\frac{\varphi(a)+\varphi(b)}{2}\right)\right|^{q}+\left|f'\left(\varphi(b)\right)\right|^{q}\right)^{\frac{1}{q}} \end{array}\right]
\end{aligned} \tag{2.7}$$

3. *For $\varphi(t)=t$ in (2.7) inequality :*

$$\begin{aligned}
&\left|\frac{f\left(a\right)+f\left(b\right)}{2}-\frac{1}{b-a}\int_{a}^{b} f(x)dx\right| \\
\leq\ & \frac{b-a}{4}\left(\frac{1}{(p+1)}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1}{q}}\left[\left(\left|f'\left(\frac{a+b}{2}\right)\right|^{q}+\left|f'\left(a\right)\right|^{q}\right)^{\frac{1}{q}}+\left(\left|f'\left(\frac{a+b}{2}\right)\right|^{q}+\left|f'\left(b\right)\right|^{q}\right)^{\frac{1}{q}}\right]
\end{aligned}$$

   we get the same result in [5] *for $s=1$.*

(4) *If we take* $\varphi(t) = t$ *in (2.6) for strongly convex functions with modulus* $c > 0$ *on* $[a,b]$ *inequality, then we obtain*

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)dx\right| \\
\leq\ & \frac{b-a}{4}\left(\frac{1}{(p+1)}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1}{q}}\left[\begin{array}{c} \left(\left|f'\left(\frac{a+b}{2}\right)\right|^q + |f'(a)|^q - \frac{c}{3}(b-a)^2\right)^{\frac{1}{q}} \\ +\left(\left|f'\left(\frac{a+b}{2}\right)\right|^q + |f'(b)|^q - \frac{c}{3}(b-a)^2\right)^{\frac{1}{q}} \end{array}\right]
\end{aligned}$$

Theorem 4. *Let* $f : I \subset \mathbb{R} \to \mathbb{R}$ *be a differentiable mapping on* I°, $a, b \in I$ *with* $a < b$ *such that* $f' \in L[a,b]$. *If* $|f'|^q$ *strongly* $\varphi-$*convex functions with modulus* c *on* $[a,b]$ *for the continuous function* $\varphi : [a,b] \longrightarrow [a,b]$ *with* $\varphi(a) < \varphi(b)$ *and* $q > 1$ *then*

$$\begin{aligned}
&\left|\frac{f(\varphi(a))+f(\varphi(b))}{2} - \frac{1}{\varphi(b)-\varphi(a)}\int_{\varphi(a)}^{\varphi(b)} f(x)dx\right| \\
\leq\ & \frac{\varphi(b)-\varphi(a)}{2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{|f'(\varphi(b))|^q + |f'(\varphi(a))|^q}{2} - \frac{c}{6}(\varphi(b)-\varphi(a))^2\right)^{\frac{1}{q}}.
\end{aligned} \tag{2.8}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From Lemma 1 and using the Hölder's integral inequality, we have

$$\begin{aligned}
&\left|\frac{f(\varphi(a))+f(\varphi(b))}{2} - \frac{1}{\varphi(b)-\varphi(a)}\int_{\varphi(a)}^{\varphi(b)} f(x)dx\right| \\
\leq\ & \frac{\varphi(b)-\varphi(a)}{2}\int_0^1 |2t-1|\,|f'(t\varphi(b)+(1-t)\varphi(a))|\,dt \\
\leq\ & \frac{\varphi(b)-\varphi(a)}{2}\left(\int_0^1 |2t-1|^p\,dt\right)^{\frac{1}{p}}\left(\int_0^1 |f'(t\varphi(b)+(1-t)\varphi(a))|^q\,dt\right)^{\frac{1}{q}} \\
\leq\ & \frac{\varphi(b)-\varphi(a)}{2}\left(\frac{1}{(p+1)}\right)^{\frac{1}{p}}\left(\int_0^1 t\,|f'(\varphi(b))|^q + (1-t)\,|f'(\varphi(a))|^q - ct\,(1-t)\,(\varphi(b)-\varphi(a))^2\,dt\right)^{\frac{1}{q}} \\
=\ & \frac{\varphi(b)-\varphi(a)}{2}\left(\frac{1}{(p+1)}\right)^{\frac{1}{p}}\left(\frac{|f'(\varphi(b))|^q + |f'(\varphi(a))|^q}{2} - \frac{c}{6}(\varphi(b)-\varphi(a))^2\right)^{\frac{1}{q}}.
\end{aligned}$$

□

Corollary 3. *Suppose that all the assumptions of Theorem4 are satisfied, in this case:*

(1) For $c = 0$, i.e. if $|f'|^q$ is $\varphi-$convex functions, we have

$$\left|\frac{f(\varphi(a)) + f(\varphi(b))}{2} - \frac{1}{\varphi(b) - \varphi(a)} \int\limits_{\varphi(a)}^{\varphi(b)} f(x)dx\right| \leq \frac{\varphi(b) - \varphi(a)}{2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{|f'(\varphi(b))|^q + |f'(\varphi(a))|^q}{2}\right)^{\frac{1}{q}} \tag{2.9}$$

(2) We obtained the same result in [3, Theorem 2.3] for $\varphi(t) = t$ in (2.9) inequality :

$$\left|\frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int\limits_{a}^{b} f(x)dx\right| \leq \frac{b-a}{2}\left(\frac{1}{(p+1)}\right)^{\frac{1}{p}}\left(\frac{|f'(a)|^q + |f'(b)|^q}{2}\right)^{\frac{1}{q}}$$

(3) If we take $\varphi(t) = t$ in (2.8) for strongly convex functions with modulus $c > 0$ on $[a, b]$ inequality, then we obtain

$$\left|\frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int\limits_{a}^{b} f(x)dx\right| \leq \frac{b-a}{4}\left(\frac{1}{(p+1)}\right)^{\frac{1}{p}}\left(\frac{|f'(a)|^q + |f'(b)|^q}{2} - \frac{c}{6}\left(\varphi(b) - \varphi(a)\right)^2\right)^{\frac{1}{q}}$$

References

[1] M. Bessenyei and Zs. Páles, *Hadamard-type inequalities for generalized convex functions*, Math. Inequal. Appl. 6/3, 279-392, 2003.

[2] M.K. Bakula, J. Pečarić, *Note on some Hadamard type inequalities*, Journal of Inequalities in Pure and Applied Mathematics, vol. 5, article 74, 2004.

[3] S.S.Dragomir,R.P.Agarwal,*Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula,* Applied Mathematics Letters, 11 (5) (1998) 91–95.

[4] S.S. Dragomir and C.E.M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications,* RGMIA Monographs, Victoria University, 2000.

[5] U.S. Kirmaci, M.K. Bakula, M.E. Özdemir and J. Pečarić, *Hadamard-type inequalities for s-convex functions,*Applied Mathematics and Computation 193 (2007) 26–35.

[6] N. Merentes and K. Nikodem, *Remarks on strongly convex functions,* Aequationes Math. 80 (2010), no. 1-2, 193-199.

[7] K. Nikodem and Zs. Pales, *Characterizations of inner product spaces be strongly convex functions,* Banach J. Math. Anal. 5 (2011), no. 1, 83-87.

[8] C.E.M. Pearce, J. Pečarić, *Inequalities for differentiable mappings with application to special means and quadrature formulæ,* Applied Mathematics Letters, 13 (200), 51-55.

[9] B.T. Polyak, *Existence theorems and convergence of minimizing sequences in extremum problems with restictions,* Soviet Math Dokl. 7 (1966), 72-75.

[10] M.Z. Sarikaya, *On Hermite Hadamard-type inequalities for strongly $\varphi-$convex functions*, Studia Universitatis Babes-Bolyai Mathematica, in press.

[11] E.Set, M.E. Özdemir and S.S. Dragomir, *On Hadamard-Type inequalities involving several kinds of convexity*, Journal of Inequalities and Applications, Article ID 286845, 12 pages, 2010.

[12] E.A. Youness, *E-convex sets, E-convex functions and E-convex programming,*Journal of Optimization Theory and Applications, 102, 2 (1999), 439-450.

Department of Mathematics, Faculty of Arts and Sciences,, Giresun University, 28100, Giresun, Turkey.
E-mail address: `imdat.iscan@giresun.edu.tr; imdati@yahoo.com`

Department of Mathematics, Faculty of Arts and Sciences,, Ordu University, 52200, Ordu, Turkey.
E-mail address: `erdalunluyol@odu.edu.tr; eunluyol@yahoo.com`